MIECHV BENCHMARKS: PM 4


WELL CHILD VISITS: FY 2020-2023

PERFORMANCE MEASURE (PM) 4 DEFINITION

Percent of TCs (target children) who received the LAST recommended well child visit in the reporting period based on the American Academy of Pediatrics (AAP) schedule

ALL HFNY MIECHV PERFORMANCE


• About 50% of TCs across all MIECHV programs received the last recommended well child visit in the reporting period over the last 3.5 years



DISPARITIES IN WELL CHILD VISITS (FY 2022)

Correlates of TC Receiving Last Well Child Visit

TCs with foreign-born or Hispanic PC1s were much more likely to have received their last well child visit than TCs of non-Hispanic White PC1s*



*Groups that were not significantly different from the reference group (non-Hispanic White PC1s) are not shown.







Other Key Findings

TCs who had health insurance were more than twice as likely to receive their last well child visit.



TCs in families with adequate food, clothing, and household goods (as reported by PC1s at intake or their last follow-up visit) were 60% more likely to receive their last well child visit.

MIECHV BENCHMARKS: PM 4

WELL CHILD VISITS: FY 2020-2023

WELL CHILD VISITS & IMMUNIZATIONS (FY 2022)												
When TCs had immunizations due, they were much more likely to receive their well child visit.												
Well Child Visit Interval	3-7 days	2 wks. - 1 mo.	2-3 mos.	4-5 mos.	6-7 mos.	9-10 mos.	12-13 mos.	15-16 mos.	18-19 mos.	24-30 mos.	36-42 mos.	48-54 mos.
% Achieved	44%	35%	80%	83%	82%	34%	81%	56%	65%	65%	49%	56%
Vaccine Interval	Birth	1 mo.	2 mos.	4 mos.	6 mos.	9 mos.	12 mos.	15 mos.	18 mos.	2-3	yrs.	4-6 yrs.
Hepatitis B	1st dose	2nd	dose				3rd dose					
Rotavirus			1st dose	2nd dose						-		
Diphtheria, tetanus & acellular pertussis			1st dose	2nd dose	3rd dose			4th	4th dose			5th dose
Haemophilus influenzae type b			1st dose	2nd dose			3rd or 4	th dose				
Pneumococcal conjugate			1st dose	2nd dose	3rd dose		3rd or 4	th dose				
Inactivated poliovirus			1st dose	2nd dose			3rd dose		1			4th dose
COVID-19							2- or 3-dose primary series and booster					
Influenza: IIV4 OR LAIV4					Annua	al vaccinat	ion (1 or 2 d	doses)	Annual va	Annual	vaccinatio doses)	n (1 or 2
Measles, mumps, rubella							1st c	dose				2nd dose
Varicella							1st c	dose				2nd dose
Hepatitis A							2	-dose seri	es			
Hepatitis A							2	-dose seri	es			







This infographic was developed by the Center for Human Services Research under an agreement with the New York State Office of Children and Family Services.